

Europäisches Patentamt
European Patent Office
Office européen des brevets



(11) **EP 1 219 613 A1**

(12)

EUROPEAN PATENT APPLICATION

(43) Date of publication:
03.07.2002 Bulletin 2002/27

(51) Int Cl.⁷: **C07D 277/32**

(21) Application number: **01130397.1**

(22) Date of filing: **20.12.2001**

(84) Designated Contracting States:
AT BE CH CY DE DK ES FI FR GB GR IE IT LI LU
MC NL PT SE TR
Designated Extension States:
AL LT LV MK RO SI

(30) Priority: **28.12.2000 JP 2000400802**

(71) Applicant: **Takeda Chemical Industries, Ltd.**
Osaka-shi, Osaka 541 (JP)

(72) Inventors:
• **Miyazaki, Takashi**
Hikari-shi, Yamaguchi (JP)

• **Satou, Makoto**
Hikari-shi, Yamaguchi (JP)
• **Inoue, Yoshihisa**
Hikari-shi, Yamaguchi (JP)

(74) Representative:
von Kreisler, Alek, Dipl.-Chem. et al
Patentanwälte,
von Kreisler-Selting-Werner,
Bahnhofsvorplatz (Deichmannhaus)
50667 Köln (DE)

(54) **Process for preparing 2-chloro-5-chloromethylthiazole**

(57) A novel process for preparing 2-chloro-5-chloromethylthiazole is provided which is suitable for industrial application.

The process for preparing 2-chloro-5-chloromethylthiazole involves allowing 2-halogenoallyl isothiocy-

anate to react with chlorinating agent in the presence of an aromatic hydrocarbon which may have one or more substituents.

EP 1 219 613 A1

Description

BACKGROUND OF THE INVENTION

5 Field of the Invention

[0001] The present invention relates to a novel process for preparing 2-chloro-5-chloromethylthiazole.

Description of the Prior Art

10

[0002] 2-chloro-5-chloromethylthiazole is an important compound as an intermediate product in a process for preparing biologically active compounds such as insecticides (see, for example, Japanese Patent Application Laid-Open No. 3-157308 (1991)). It can be prepared by, for example, the following processes.

15

[0003] Japanese Patent Application Laid-Open No. 63-83079 (1988) discloses a process for preparing 2-chloro-5-chloromethylthiazole by allowing allyl isothiocyanate to react with chlorinating agent as described in the following scheme (A):

20



25

(A)


[0004] This process requires a large amount of chlorinating agent and the reaction has to be performed under a high temperature, which produces many by-products.

30

Therefore, it is not so suitable for industrial application.

[0005] Alternatively, Japanese Patent Application Laid-Open No. 4-234864 discloses a process for preparing 2-chloro-5-chloromethylthiazole by allowing allyl isothiocyanate derivative to react with chlorinating agent as described in the following scheme (B):

35



40

X=leaving group

(B)

45

[0006] This process is suitable for industrial application which can be performed using approximately an equal amount of chlorinating agent under moderate condition to give a high yield. However, inert solvent (diluent) such as halocarbon may preferably be used to give a high yield, which requires high production cost.

50

[0007] Japanese Patent Application Laid-Open No. 2000-247963 (2000) discloses a process which involves allowing 2-halogenoallyl isothiocyanate to react with chlorinating agent in a dipolar aprotic solvent such as acetonitrile. This process, however, has a disadvantage that the solvent cannot be reused since it is difficult to separate and collect the solvent such as acetonitrile from water.

55

[0008] Accordingly, a novel process for preparing 2-chloro-5-chloromethylthiazole suitable for industrial application is still needed in which an excess amount of chlorinating agent is not required, moderate conditions can be used to give a high yield, and solvent can be collected easily.


[0009] The object of the present invention to provide a novel process for preparing 2-chloro-5-chloromethylthiazole which is suitable for industrial application.

SUMMARY OF THE INVENTION


[0010] Aromatic hydrocarbons and derivatives thereof (particularly toluene) are reactive with chlorinating agent. The present inventors unexpectedly found, as a result of intense studies, that 2-chloro-5-chloromethylthiazole can be obtained at a high yield by allowing 2-halogeno-allyl isothiocyanate to react with chlorinating agent in the presence of aromatic hydrocarbons which may have one or more substituents, and accomplished the present invention based on the findings.

[0011] In other words, the present invention provides:

[0012] (1) a process for preparing 2-chloro-5-chloromethylthiazole of the following formula (I):



by allowing 2-halogeno-allyl isothiocyanate of the following formula (II):



(wherein Hal represents chlorine or bromine) to react with chlorinating agent in the presence of an aromatic hydrocarbons which may have one or more substituents;


[0013] (2) the process according to (1) above wherein the aromatic hydrocarbons which may have one or more substituents is toluene, chlorobenzene or dichlorobenzene; and

[0014] (3) the process according to (1) above wherein the aromatic hydrocarbons which may have one or more substituents is toluene.

[0015] (4) the process according to (1) above wherein Hal in the formula (II) is chlorine.

DETAILED DESCRIPTION OF THE INVENTION

[0016] A 2-halogenoallyl isothiocyanate of the following formula (II):



(wherein Hal represents chlorine or bromine) is a known compound which can be prepared by any known processes. One unlimiting example of such process involves heating a mixture of 2,3-dihalogeno-1-propene (e.g., 2,3-dichloro-1-propene) with thiocyanate represented by $M(SCN)_n$ (wherein M represents metal or ammonium group, and n indicates the valence of M) in the presence of water (see, Japanese Patent Application Laid-Open No. 9-136874 (1997)).


[0017] The process according to the present invention may be performed in the presence of an "aromatic hydrocarbons which may have one or more substituents". The aromatic hydrocarbon in the "aromatic hydrocarbon which may have one or more substituents" herein may include, for example, benzene, naphthalene and phenanthrene. The "aromatic hydrocarbon" may be substituted with, for example, C_{1-4} alkyl group (e.g., methyl, ethyl, propyl, isopropyl, butyl, isobutyl, s-butyl or t-butyl) and/or halogen (e.g., fluorine, chlorine, bromine or iodine) and number of substituents is

preferably 1 to 3. Examples of such "aromatic hydrocarbons which may have one or more substituents" are toluene, o-, m-, p-xylene and xylene isomeric mixtures, ethylbenzene, cumene, cymene, mesitylene, chlorobenzene, as well as o-, m- or p-dichlorobenzene and isomeric mixtures of dichlorobenzene. Preferable examples of "aromatic hydrocarbons which may have one or more substituents" are chlorobenzene, dichlorobenzene (o-, m- or p-dichlorobenzene or mixtures thereof) and toluene, among them toluene is particularly preferable. The "aromatic hydrocarbons which may have one or more substituents" may be used alone or in combination of two or more thereof.

[0018] Preferably 0.1 to 20 parts and more preferably 0.5 to 5 parts by weight of the "aromatic hydrocarbons which may have one or more substituents" may be used relative to 1 part by weight of the above-described 2-halogenoallyl isothiocyanate.

[0019] The term "chlorinating agent" herein refers to chlorine or a compound which releases chlorine under reaction conditions, such as sulfuryl chloride or phosgene. The chlorinating agent is typically used in an amount of 0.8 to 2 equivalents, preferably 1.0 to 1.5 equivalents, and more preferably 1.05 to 1.30 equivalents relative to the above-described 2-halogenoallyl isothiocyanate.

[0020] For example, a reaction process according to the present invention using 2-chloroallyl isothiocyanate as the starting material and sulfuryl chloride as the chlorinating agent can be particularly described with reference to the following scheme.



[0021] The process according to the present invention can be performed, for example, under the following conditions.

[0022] The reaction may typically be performed at -60 to 60°C, preferably at -10 to 50°C and more preferably at 10 to 40°C though a wide range of temperature may be selected.

[0023] The reaction may preferably be performed under atmospheric pressure though reduced, atmospheric or pressurized pressure may be used.

[0024] The reaction may typically be performed for from 10 minutes to 50 hours and preferably for 1 to 20 hours.

[0025] After reaction completed, 2-chloro-5-chloromethylthiazole can be isolated by any known methods. For example, reaction mixture may be washed with a suitable amount of water, oil layer may be separated and collected, and then the "aromatic hydrocarbons which may have one or more substituents" may be removed by, for example, vacuum distillation. The "aromatic hydrocarbons which may have one or more substituents" may be collected for recycling.

[0026] Hereinafter, the present invention will be described in more detail with reference to the following examples though they are not intended to limit the scope of the present invention.

Example 1

[0027] 2-chloroallyl isothiocyanate (75 g) was mixed with chlorobenzene (136 mL), heated to 30°C and added dropwise with sulfuryl chloride (84 g) for 3 hours. After stirring at 30°C for 2 hours, the reaction mixture was added to water (94 g). Separated chlorobenzene layer was added with water (18 g) followed by aqueous solution of 38 % potassium carbonate (48 g), and stirred for 30 minutes. Thereafter, oil layer was separated from aqueous layer. The oil layer was distilled under reduced pressure to collect chlorobenzene. The residue was further distilled under reduced pressure to give 77g of 2-chloro-5-chloromethylthiazole (purity=80 %, yield=75 %).

Example 2

[0028] 2-chloroallyl isothiocyanate (86 g) was mixed with chlorobenzene (136 mL), heated to 30°C and added dropwise with sulfuryl chloride (83 g) for 3 hours. After stirring at room temperature for 15 hours, the reaction mixture was added to water (94 g). Additionally, chlorobenzene (40mL) was added to the mixture which was then heated to 40°C and stirred for 1 hour. Thereafter, separated chlorobenzene layer was added with water (18 g) followed by dropwise aqueous solution of 38 % potassium carbonate (75 g) and stirred for 30 minutes, and oil phase was then separated from aqueous layer. The oil layer was enriched under reduced pressure to collect chlorobenzene. The enriched residue was further distilled under reduced pressure to give 75 g of 2-chloro-5-chloromethylthiazole (purity=95 %, yield=76 %).

•Example 3

[0029] 2-chloroallyl isothiocyanate (30 g) was mixed with toluene (40 mL), and added dropwise with sulfuryl chloride (45 g) at room temperature for 1 hour. After stirring at room temperature for 2 hours, deposited crystal was collected by filtration, washed with toluene (40 mL), mixed with water (40 mL), heated to 40°C and then cooled to room temperature, and added with aqueous solution of 38 % potassium carbonate (16 g) to adjust pH to approximately 2. Then, oil layer was separated from aqueous layer. The oil layer was distilled under reduced pressure to give 26.0 g of 2-chloro-5-chloromethylthiazole (purity=95 %, yield=73 %).

10 Example 4

[0030] 2-chloroallyl isothiocyanate (30 g) was mixed with toluene (40 mL), and added dropwise with sulfuryl chloride (45 g) at room temperature for 1 hour. After stirring at room temperature for 2 hours, the mixture was heated to 40°C, stirred for 1 hour, and then added dropwise with water (5 mL). Then, aqueous solution of 38 % potassium carbonate (16 g) was added at room temperature to adjust pH to approximately 2. Then, water (10 mL) was added and separated. Toluene layer was enriched under reduced pressure to collect toluene (80 %) and then the enriched residue was further distilled under reduced pressure to give 27.1 g of 2-chloro-5-chloromethylthiazole (purity=98.0 %, yield=79 %).

Example 5

[0031] 2-chloroallyl isothiocyanate (purity=95.5%, 94.2 kg) was mixed with chlorobenzene (181.8 kg), heated to 30°C and added dropwise with sulfuryl chloride (101.3 kg) for 4 hours. After stirring at 30°C overnight, the reaction mixture was added to water (113.6 kg). Separated chlorobenzene layer was added with water (22.2 kg) followed by aqueous solution of 38 % potassium carbonate (39.1 kg), stirred for 1 hour, and allowed to stand overnight. Thereafter, organic layer was separated from aqueous layer. The organic layer was distilled under reduced pressure to collect chlorobenzene. The residue was further distilled under reduced pressure (102-104°C / 16-14 mmHg) to give 87.4 kg of 2-chloro-5-chloromethylthiazole (purity=97.4 %, yield=75.2 %).


Example 6

[0032] 2-chloroallyl isothiocyanate (purity=94.3%, 1740 kg) was mixed with toluene (1958 kg), heated to 30°C and added dropwise with sulfuryl chloride (1892 kg) for 4 hours. After stirring at 40°C for 2 hours, the mixture was added with water (1538 kg) followed by aqueous solution of 38 % potassium carbonate (2393 kg), and stirred for 30 minutes, and oil layer was then separated from aqueous layer. The oil layer was enriched under reduced pressure to collect toluene. The enriched residue was further distilled under reduced pressure (86-90°C / 4-2 mmHg) to give 1692 kg of 2-chloro-5-chloromethylthiazole (purity=95.0 %, yield=77.9 %).


[0033] As described above, a novel process for preparing 2-chloro-5-chloromethylthiazole is provided according to the present invention in which an excess amount of chlorinating agent is not required, moderate conditions can be used to give a high yield, and solvent can be collected easily after reaction completed.

Claims

1. A process for preparing 2-chloro-5-chloromethylthiazole of the following formula (I):



by allowing 2-halogeno-allyl isothiocyanate of the following formula (II):



(wherein Hal represents chlorine or bromine) to react with chlorinating agent in the presence of an aromatic hydrocarbon which may have one or more substituents.

2. The process according to claim 1 wherein the aromatic hydrocarbon which may have one or more substituents is toluene, chlorobenzene or dichlorobenzene.
3. The process according to claim 1 wherein the aromatic hydrocarbon which may have one or more substituents is toluene.
4. The process according to claim 1 wherein Hal in the formula (II) is chlorine.



European Patent
Office

EUROPEAN SEARCH REPORT

Application Number
EP 01 13 0397

DOCUMENTS CONSIDERED TO BE RELEVANT			
Category	Citation of document with indication, where appropriate, of relevant passages	Relevant to claim	CLASSIFICATION OF THE APPLICATION (Int.Cl.7)
Y	EP 0 794 180 A (KURARAY CO., LTD., JAPAN) 10 September 1997 (1997-09-10) * column 7, line 50 - column 8, line 13 *	1-4	C07D277/32
Y	WO 97 10226 A (CIBA GEIGY AG ;PITTERNA THOMAS (CH)) 20 March 1997 (1997-03-20) variant b2	1-4	
Y,D	EP 0 446 913 A (TAKEDA CHEMICAL INDUSTRIES LTD) 18 September 1991 (1991-09-18) * the whole document *	1-4	
A,D	EP 1 031 566 A (BAYER AG) 30 August 2000 (2000-08-30) * the whole document *	1-4	
A,D	EP 0 260 560 A (BAYER AG) 23 March 1988 (1988-03-23) * the whole document *	1-4	
The present search report has been drawn up for all claims			TECHNICAL FIELDS SEARCHED (Int.Cl.7)
			C07D
Place of search MUNICH		Date of completion of the search 20 February 2002	Examiner Bérillon, L
<p>CATEGORY OF CITED DOCUMENTS</p> <p>X : particularly relevant if taken alone Y : particularly relevant if combined with another document of the same category A : technological background O : non-written disclosure P : intermediate document</p> <p>T : theory or principle underlying the invention E : earlier patent document, but published on, or after the filing date D : document cited in the application L : document cited for other reasons & : member of the same patent family, corresponding document</p>			

EPO FORM 1503 03 02 (P4001)

**ANNEX TO THE EUROPEAN SEARCH REPORT
ON EUROPEAN PATENT APPLICATION NO.**

EP 01 13 0397

This annex lists the patent family members relating to the patent documents cited in the above-mentioned European search report. The members are as contained in the European Patent Office EDP file on
The European Patent Office is in no way liable for these particulars which are merely given for the purpose of information.

20-02-2002

Patent document cited in search report		Publication date		Patent family member(s)	Publication date
EP 0794180	A	10-09-1997	JP	9316062 A	09-12-1997
			JP	10036342 A	10-02-1998
			JP	10045709 A	17-02-1998
			JP	10045710 A	17-02-1998
			AU	690866 B2	30-04-1998
			AU	1481997 A	28-08-1997
			CN	1173496 A ,B	18-02-1998
			EP	0794180 A1	10-09-1997
			JP	10036360 A	10-02-1998
			US	6103921 A	15-08-2000
			US	6222057 B1	24-04-2001
			US	6245927 B1	12-06-2001
			US	5894073 A	13-04-1999
WO 9710226	A	20-03-1997	AU	7084196 A	01-04-1997
			WO	9710226 A1	20-03-1997
			EP	0850231 A1	01-07-1998
			JP	11512407 T	26-10-1999
			US	6187927 B1	13-02-2001
EP 0446913	A	18-09-1991	AT	154016 T	15-06-1997
			CA	2038203 A1	17-09-1991
			DE	69126350 D1	10-07-1997
			DE	69126350 T2	06-11-1997
			DK	446913 T3	07-07-1997
			EP	0446913 A1	18-09-1991
			ES	2103753 T3	01-10-1997
			GR	3024285 T3	31-10-1997
			HU	59675 A2	29-06-1992
			IL	97565 A	29-06-1995
			JP	3179515 B2	25-06-2001
			JP	4234864 A	24-08-1992
			KR	215132 B1	16-08-1999
			US	5180833 A	19-01-1993
EP 1031566	A	30-08-2000	DE	19908447 A1	31-08-2000
			BR	0000962 A	19-09-2000
			CN	1266848 A	20-09-2000
			EP	1031566 A1	30-08-2000
			JP	2000247963 A	12-09-2000
			US	6214998 B1	10-04-2001
EP 0260560	A	23-03-1988	DE	3631538 A1	24-03-1988
			CA	1306463 A1	18-08-1992
			DD	263988 A5	18-01-1989
			DE	3762745 D1	21-06-1990

C/O FORM P0439

For more details about this annex : see Official Journal of the European Patent Office, No. 12/82

**ANNEX TO THE EUROPEAN SEARCH REPORT
ON EUROPEAN PATENT APPLICATION NO.**

EP 01 13 0397

This annex lists the patent family members relating to the patent documents cited in the above-mentioned European search report. The members are as contained in the European Patent Office EDP file on
The European Patent Office is in no way liable for these particulars which are merely given for the purpose of information.

20-02-2002

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
EP 0260560	A	EP 0260560 A1	23-03-1988
		HU 48225 A2	29-05-1989
		JP 2078542 C	09-08-1996
		JP 7116167 B	13-12-1995
		JP 63083079 A	13-04-1988
		KR 9411458 B1	15-12-1994
		US 4748243 A	31-05-1988

EPO FORM P0159

For more details about this annex : see Official Journal of the European Patent Office, No. 12/82